


Yes

You've been identified as a close contact of someone who has tested positive for COVID-19 or someone with COVID-19 symptoms. Now what?

This guidance does not apply to individuals who live, work, volunteer or are admitted in a highest risk setting (hospitals, Long-Term Care, retirement homes, congregate living settings)



- at least 10 days (or 5 days if under 12 years old) following your last exposure $^{\Omega}$, unless you have previously tested positive for COVID-19 in the last 90 days.
- If you are eligible for testing, get tested and follow the auidance for cases.

- Report your exposure to your employer and follow any work restrictions.
- If you develop symptoms, self-isolate immediately for at least 5 days from symptom onset and until you have no fever and other symptoms are improving for 24 hours (or 48 hours for gastrointestinal symptoms).
- Get tested if eligible and follow the guidance for cases.

- Self-isolate for the same amount of time as the positive/symptomatic person.**
 - If you have previously tested positive for COVID-19 in the last 90 days and are asymptomatic, you do not need to isolate.
- If you develop <u>symptoms</u> continue/start to self-isolate and get tested if you are eligible. Follow the guidance for cases.
- If anyone else in the household develops symptoms then all other asymptomatic household contacts should extend their self-isolation until the newly symptomatic person has finished their self-isolation period.

Wear a well-fitted mask in public, physical distance and maintain other public health measures for 10 days following your last exposure if leaving home. You should **NOT visit any highest risk settings or individuals who may be at higher risk of illness (i.e. seniors or immunocompromised) for 10 days from your last exposure. Immunocompromised individuals may still be susceptible regardless of vaccination and should self-isolate for 10 days after last exposure to a case/symptomatic person.

 $^{\Omega}$ If you develop <u>symptoms</u>, <u>self-isolate</u> for at least 10 days (or 5 days if under 12 years old) from symptom onset and until you have no fever and other symptoms are improving for 24 hours (or 48 hours for gastrointestinal symptoms).